

NEWS

America's Frontline Doctors: COVID-vaccinated can 'shed' spike protein, harming unvaccinated

As these experimental vaccines create 'spike proteins,' vaccinated individuals 'can shed some of these particles to close contacts' causing disease in them, including in children.

Mon May 3, 2021 - 1:14 pm EST



SHUTTERSTOCK.COM

By Patrick Delaney

LOS ANGELES, California, May 3, 2021 (LifeSiteNews) — In their latest issue brief, America's Frontline Doctors (AFLDS) warned how spike proteins resulting from experimental COVID-19 gene therapy vaccines have the capacity to 1.) pass through the "blood-brain barrier" causing neurological damage, 2.) be "shed" by the vaccinated,

bringing about sickness in unvaccinated children and adults, and 3.) cause irregular vaginal bleeding in women.

Released last week and titled “Identifying Post-vaccination Complications & Their Causes: an Analysis of Covid-19 Patient Data,” the stated purpose of the document is “to provide additional information for concerned citizens, health experts, and policymakers about adverse events and other post-vaccination issues resulting from the three experimental COVID-19 vaccines currently administered under EUA (emergency use authorization)” by the U.S. Food and Drug Administration (FDA).

The non-profit organization highlighted the thousands of adverse events which are related to these “vaccines” and captured by the Centers for Disease Control and Prevention’s (CDC) Vaccine Adverse Event Reporting System (VAERS). “Yet these complications have received a fraction of the attention paid to J&J’s blood-clotting controversy,” they lamented with dismay, asking, “Why?”

In taking a closer look at this data, AFLDS presents “some major categories of concern as-yet publicly unaddressed by either the FDA or CDC,” asserting that failure of these regulators “to consider these and other ‘known unknowns’ is a dereliction of basic medical research.”

They breakout their general categories of concern as shown below:

First, there are significant fears regarding the wide distribution of these new vaccines, which employ a new technology and remain only experimental without full approval from the FDA. Instead of employing an attenuated antigen response – as happens with conventional vaccines – these experimental agents introduce something called a “spike protein” into one’s system.

“It takes years to be sure something new is safe,” the AFLDS document confirms. “No one knows definitively the long-term health implications for the body and brain, especially among the young, related to this spike protein. In addition, if documented problems with the protein do arise, there will never be any way to reverse the adverse effects in those already vaccinated.”

Second, unlike conventional vaccines, these spike proteins, along with “lipid nanoparticles” have the capacity to pass through the “blood-brain barrier” which provides special protection for these sensitive areas of the body.

“There simply has not been enough time to know what brain problems and how often a brain problem will develop from that,” the document warns.

Risks from such penetration include “chronic inflammation and thrombosis (clotting) in the neurological system, contributing to tremors, chronic lethargy, stroke, Bell’s Palsy and ALS-type symptoms. The lipid nanoparticles can potentially fuse with brain cells, resulting in delayed neuro-degenerative disease. And the mRNA-induced spike protein can bind to

brain tissue 10 to 20 times stronger than the spike proteins that are (naturally) part of the original virus.”

Third, as these experimental vaccines produce many trillions of spike proteins in their recipients, these vaccinated individuals “can shed some of these (spike protein) particles to close contacts,” causing disease in them.

In an email correspondence with LifeSiteNews, Dr. Simone Gold, the founder of AFLDS, directed this writer to an April 29 tweet where she posted a document from Pfizer’s experimental trials in which the pharmaceutical giant “acknowledges this mechanism” of potential shedding, she wrote.

Contradictory much? Pfizer trials warned men to stay away from pregnant women... but now the CDC is pushing pregnant women to take an experimental biological agent without a second thought. pic.twitter.com/rJkeDNFRvE

— Dr. Simone Gold (@drsimonegold) April 29, 2021

As the document states, one can be “exposed to [the] study intervention due to environmental exposure,” including “by inhalation or skin contact” with someone involved in the study, or with another who has been exposed in the same way.

And this, according to AFLDS, can be dangerous. As the issues brief continues, “the spike proteins are pathogenic (‘disease causing’) just like the full virus.” Furthermore, these “spike proteins bind *more* tightly than the fully intact virus” and thus cases around the world of “pericarditis, shingles, pneumonia, blood clots in the extremities and brain, Bell’s Palsy, vaginal bleeding and miscarriages have been reported in persons who are near persons who have been vaccinated.” Such shedding also “appears to be causing wide variety of autoimmune disease (where the body attacks its own tissue) in some persons.”

— Article continues below Petition —

In addition, other more serious dangers to even the unvaccinated are possible due to the fact that these “spike proteins can cross the blood brain barrier, unlike traditional vaccines.”

Fourth, such shedding leaves children vulnerable if they are in proximity to parents and teachers who have received these experimental vaccines. While the threat of COVID-19 to the young is rightly described as “irrelevant,” including a 99.997% survival rate for those under 20 years of age, AFLDS is concerned some children may become symptomatic due to such proximity to the vaccinated. At such point there is a danger that “public health bureaucrats” might use such cases to “speculate that a child’s illness is related to a SARS-CoV-2 ‘variant,’” when it is a result of contact with vaccinated adults.

“Our other concern is that children could develop long-term chronic autoimmune disease including neurological problems due to the fact that children have decades ahead of them

including neurological problems due to the fact that children have decades ahead of them and trillions of the spike proteins mentioned above.”

Fifth, “AFLDS is aware of thousands of reports involving vaginal bleeding, post-menopausal vaginal bleeding, and miscarriages following COVID-19 vaccination as well as anecdotal reports of similar adverse events among those in close contact with the vaccinated.” While at this point the independent physicians organization “cannot comment definitively on the close contacts” other than to mention they “have heard reports of this worldwide,” the many reported incidents of post-vaccination vaginal bleeding establishes a clear “connection between the vaccine and irregular bleeding.”

“Despite this clear-cut evidence, menstrual-cycle changes were not listed among the FDA’s common side effects in its phase-three clinical participants. Women’s reproductive health needs to be taken seriously rather than waved away by agenda-driven public health officials,” the brief reads.

Finally, acknowledging the “irrepressible economic incentive among pharmaceutical companies” to market unnecessary and dangerous childhood COVID vaccines, boosters, and the like, AFLDS insists “Public health experts should stop and assess data on possible vaccine side effects and related post-vaccination questions before it is too late.”

RELATED:

Thousands of reports of menstrual irregularities, reproductive dysfunction following COVID vaccines

Frontline Doctors: Experimental vaccines are ‘not safer’ than COVID-19

Dr. Peter McCullough: given an 80% level of herd immunity, broad vaccination has ‘no scientific, clinical or safety rationale.’

EXCLUSIVE - Former Pfizer VP: ‘Your government is lying to you in a way that could lead to your death.’

Anti-lockdown scientists challenge theories of Geert Vanden Bossche, though vaccine ‘global catastrophe’ not ruled out

‘Miraculous’ ivermectin approved for use in the US for the treatment of COVID-19
